

The first and only FDA-approved, once-weekly treatment
for use in children with achondroplasia*

What to expect with



It's natural to have questions about your child's treatment journey.

This brochure has helpful information about:

- What to know before starting treatment with YUWIWEL
- Where you can find Instructions for Use
- How to connect with patient support

***What is YUWIWEL?**

YUWIWEL is a prescription medicine used to increase linear growth in children 2 years and older with achondroplasia with open growth plates (epiphyses).

YUWIWEL is approved under accelerated approval based on an improvement in annualized growth velocity. Continued approval may be contingent upon verification and description of clinical benefit in confirmatory trials.

IMPORTANT SAFETY INFORMATION

What are the possible side effects of YUWIWEL?

YUWIWEL may cause serious side effects, including risk of low blood pressure. If your child experiences a decrease in blood pressure or symptoms of low blood pressure (dizziness, feeling tired, and/or nausea) while being treated with YUWIWEL, call your child's healthcare provider.

Please see Important Safety Information throughout and full Prescribing Information and Patient Information for YUWIWEL.

How can YUVIWEL help my child grow?

YUVIWEL can be part of a treatment plan designed to help your child grow.




YUVIWEL is a once-weekly injection



Your child can take YUVIWEL until their growth plates close

Growth plates are the parts of bones where new growth happens. They usually close by age 15 in girls and by age 17 in boys.



IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of YUVIWEL? (cont'd)

The most common side effects of YUVIWEL include injection site reactions (redness, itching, skin discoloration, bleeding, swelling, bruising, pain, and blistering).

Please see Important Safety Information throughout and full Prescribing Information and Patient Information for YUVIWEL.

You'll have a team supporting you as you go

When your family decides to start YUVIWEL:

- The Ascendis Signature Access Program® (A-S-A-P) is available every step of the way. From starting treatment to navigating insurance, A-S-A-P can help you stay on track*
- A-S-A-P can help arrange for a Patient Access Liaison (PAL) to provide injection training support and education
- With A-S-A-P, a dedicated team is available to answer questions you may have

A · S · A · P
 ASCENDIS SIGNATURE
 ACCESS PROGRAM®



Want some help? Call 1-844-442-7236 to get in contact with an A-S-A-P team member, or scan the QR code to learn more about the program

*The A-S-A-P program helps provide patient support services to eligible patients who were prescribed YUVIWEL for its approved indication.

How YUVIWEL may help

YUVIWEL results


The YUVIWEL study was made up of children aged 2 to 12. The study measured how many inches a child grew in a year.

- This measurement is called annualized growth velocity (AGV)

In the study:

- **Children getting YUVIWEL grew faster**, on average, than those getting placebo (an injection with no medicine)
- **The YUVIWEL group grew 0.6 inches (1.5 cm) more per year** than the placebo group

Growth per year with YUVIWEL vs placebo



4 Please see Important Safety Information throughout and full Prescribing Information and Patient Information for YUVIWEL.

YUVIWEL safety profile

In studies, YUVIWEL was generally well tolerated

- The most common side effects were vomiting, injection site reactions, pain in the arms and legs, and nausea
 - Injection site reactions included redness, itching, skin discoloration, bleeding, swelling, bruising, pain, and blistering
- All injection site reactions were mild and did not cause any children to stop taking YUVIWEL
- 3% of children treated with YUVIWEL saw an increase in hair growth either near the injection site or on the limbs, back, and shoulders, compared with 0% of children who received placebo. Rotating the site of injection with each dose can reduce this risk



If you have any questions about your child's treatment with YUVIWEL, you can always reach out to your Patient Access Liaison (PAL) for support.

IMPORTANT SAFETY INFORMATION (cont'd)

Before you give YUVIWEL to your child, tell the healthcare provider about all of your child's medical conditions, including if they:

- have kidney problems
- are pregnant or plan to become pregnant. It is not known if YUVIWEL will harm the unborn baby
- are breastfeeding or plan to breastfeed. It is not known if YUVIWEL passes into breast milk

Ready to start? Here are some things to know

Dosing:



YUVIWEL can be taken with or without food



The YUVIWEL dose is based on weight. Your care team will keep track of how much your child weighs and adjust their dose as needed

Once the team confirms that their growth plates have closed, they should stop taking YUVIWEL



YUVIWEL is given once weekly



Try to give it on the same day each week

YUVIWEL can be taken up to 2 days before or 2 days after the planned dosing day*

A Patient Access Liaison (PAL) will help arrange injection training support at your home.


YUVIWEL must be carefully mixed and prepared before administering. Please refer to the Instructions for Use or ask your child's doctor for complete instructions.

*If more than 2 days have passed, skip the missed dose, and the next dose should be given on the regularly scheduled dosing day. In both cases, the regular once-weekly dosing schedule can be resumed. At least 5 days must pass between doses.

Giving your child YUVIWEL

YUVIWEL is injected under the skin. This is called a subcutaneous injection.

Where should YUVIWEL be given?



Each box of medication has everything you need for 4 YUVIWEL injections

The YUVIWEL Instructions for Use will help explain all of the items and how they're used



Store unmixed YUVIWEL in a refrigerator. YUVIWEL can also be stored at room temperature (up to 86 °F) for up to 6 months and can be returned to refrigeration within 6 months

YUVIWEL does not contain preservatives. After mixing, YUVIWEL can be stored at room temperature and must be used within 4 hours. Throw away unused medicine



Scan code to watch step-by-step instructions on how to prepare YUVIWEL

IMPORTANT SAFETY INFORMATION (cont'd)

Tell the healthcare provider about all the medicines your child takes, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Personalized support with the Ascendis Signature Access Program (A·S·A·P)



Receive one-on-one support throughout your treatment journey.

Through the Ascendis Signature Access Program (A·S·A·P), a dedicated Patient Access Liaison (PAL)* is assigned to support you with:



A direct phone number to reach them



Educational information on YUVIWEL



Information on the prior authorization process and financial assistance options[†]



Help to avoid a gap in treatment



Injection training support

*The A·S·A·P program helps provide patient support services to eligible patients who were prescribed YUVIWEL for its approved indication.

[†]Terms and Conditions apply. See Eligibility and Restrictions at yuviwel.com/terms-and-conditions.

8 Please see Important Safety Information throughout and full Prescribing Information and Patient Information for YUVIWEL.

Questions to ask your child's care team

It can be helpful to write down questions so that you can remember them for your next appointment. Here are a few to get you started.

Q: How much can I expect my child to grow?

A: _____

Q: What should I know about side effects?

A: _____

Q: How long can my child take YUVIWEL?

A: _____

Q: Can someone help me learn how to give my child YUVIWEL?

A: _____

Q: Is financial assistance available?

A: _____

Q: How do I sign up for support programs?

A: _____

IMPORTANT SAFETY INFORMATION (cont'd)

These are not all of the possible side effects of YUVIWEL. Call your doctor for medical advice about side effects. **You are encouraged to report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. You may also report side effects to Ascendis Pharma at 1-844-442-7236.**

YUWIWEL is here to help.

Your child's treatment journey will change as they grow. We offer resources and support every step of the way.

For personalized support or to learn more, call 1-844-442-7236



Scan to learn more about how YUWIWEL can help your child with achondroplasia

What is YUWIWEL?

YUWIWEL is a prescription medicine used to increase linear growth in children 2 years and older with achondroplasia with open growth plates (epiphyses).

YUWIWEL is approved under accelerated approval based on an improvement in annualized growth velocity. Continued approval may be contingent upon verification and description of clinical benefit in confirmatory trials.

IMPORTANT SAFETY INFORMATION

What are the possible side effects of YUWIWEL?

YUWIWEL may cause serious side effects, including risk of low blood pressure. If your child experiences a decrease in blood pressure or symptoms of low blood pressure (dizziness, feeling tired, and/or nausea) while being treated with YUWIWEL, call your child's healthcare provider.

Please see Important Safety Information throughout and full Prescribing Information and Patient Information for YUWIWEL.